

Ionospheric Profiles from Ultraviolet Remote Sensing

Arjun Tan

Department of Physics
Alabama A & M University
Normal, AL 35762-0447

phone: (256) 858-8115 fax: (256) 851-5622 e-mail: atan@aamu.edu

Award # N00014-97-1-0267

LONG-TERM GOALS

The long-term goal of this project is to obtain ionospheric profiles from ultraviolet remote sensing of the ionosphere from orbiting space platforms.

OBJECTIVES

Remote sensing of the nighttime ionosphere is a relatively straightforward process due to the absence of the complications brought about by daytime solar radiation. Further, during the nighttime hours, the O^+-H^+ transition level in both the mid- and low-latitude ionospheres lies around 750 km, which is within the range of accuracy of the path matrix inversion. The intensity of the O^+-e^- recombination radiation as observed from orbiting space platforms can now be used to reconstruct the nighttime ionospheric profile including the O^+-H^+ transition height.

APPROACH

In the topside ionosphere, a distinct bend in the electron density profile marks the boundary between the F region (major ion O^+) and the plasmasphere (major ion H^+) (cf. Davies, 1990). During the nighttime hours, this level is situated around 750 km for both the mid- and low-latitude ionospheres. Also, during the nighttime, chemical equilibrium condition holds for altitudes of up to at least 800 km (Hanson and Ortenburger, 1961; Geisler, 1967).

The F region density profile is approximated by a 3-parameter Chapman layer model. The nighttime O^+ density profile is first determined by direct inversion of the intensity of the O^+-e^- recombination reaction at different observation angles. The H^+ density profile is next approximated by a 3-parameter Parabolic-Chapman profile, two of which are given by the transition level altitude and density.

WORK COMPLETED

Model ionospheric profiles based on observed nighttime profiles were reconstructed using this technique. Following are some examples:

1. Lima profile (12 deg S latitude) at 1515 LT on 17 February 1962 recorded by Bowles (1963);
2. "Singapore" profile for a location over Indian Ocean (9 deg N latitude) at 2214 LT on 1 October 1962 recorded by King (1963);
3. Jicamarca profile (12 deg S latitude) at 2325 LT on 2 February 1965 recorded by McClure (1965);
and

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 1998		2. REPORT TYPE		3. DATES COVERED 00-00-1998 to 00-00-1998	
4. TITLE AND SUBTITLE Ionospheric Profiles from Ultraviolet Remote Sensing				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Alabama A&M University, Department of Physics, Normal, AL, 35762-0447				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM002252.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

4. Arecibo profile (18 deg N latitude) at 2230 LT on 10 February 1972 recorded by Hagen and Hsu (1974).

RESULTS

1. The Lima profile was approximated by an O^+ Chapman profile with $N_m = 500,000 \text{ cm}^{-3}$, $z_m = 275 \text{ km}$ and $H = 68 \text{ km}$; and by an H^+ Parabolic-Chapman profile with $N_{tr} = 2,800 \text{ cm}^{-3}$, $z_{tr} = 800 \text{ km}$ and $H = 300 \text{ km}$.
2. The “Singapore” profile was approximated by an O^+ Chapman profile with $N_m = 760,000 \text{ cm}^{-3}$, $z_m = 250 \text{ km}$ and $H = 50 \text{ km}$; and by an H^+ Parabolic-Chapman profile with $N_{tr} = 11,200 \text{ cm}^{-3}$, $z_{tr} = 800 \text{ km}$ and $H = 125 \text{ km}$.
3. The Jicamarca profile was approximated by an O^+ Chapman profile with $N_m = 105,000 \text{ cm}^{-3}$, $z_m = 426 \text{ km}$ and $H = 50 \text{ km}$; and by an H^+ Parabolic-Chapman profile with $N_{tr} = 6,500 \text{ cm}^{-3}$, $z_{tr} = 725 \text{ km}$ and $H = 155 \text{ km}$.
4. The Arecibo profile was approximated by an O^+ Chapman profile with $N_m = 20,000 \text{ cm}^{-3}$, $z_m = 350 \text{ km}$ and $H = 50 \text{ km}$; and by an H^+ Parabolic-Chapman profile with $N_{tr} = 900 \text{ cm}^{-3}$, $z_{tr} = 700 \text{ km}$ and $H = 190 \text{ km}$.

The reconstructed profiles all exhibited slightly underestimated topside O^+ densities and slightly overestimated H^+ and topside e^- densities.

IMPACT/APPLICATIONS

This study permits a more complete reconstruction of the topside ionosphere by ultraviolet remote sensing, including the lower protonosphere and the O^+ - H^+ transition altitude.

TRANSITIONS

Several forthcoming ultraviolet remote sensing experiments to be conducted by the U. S. Naval Research Laboratory will monitor the ionosphere on a continuous basis from orbiting space platforms.. The present technique can be utilized in the data analyses of these measurements.

RELATED PROJECTS

REFERENCES

- K. L. Bowles, NBS Tech. Note 169 (1963).
- K. Davies, Ionospheric Radio, Peter Peregrinus Ltd. (1990).
- J. E. Geisler, J. Geophys. Res., 72, 81 (1967).
- W. B. Hanson and I. B. Ortenburger, J. Geophys. Res., 66, 1424 (1961).
- J. W. King, Nature, 197, 639 (1963).
- J. B. Hagen and P. Y. S. Hsu, J. Geophys. Res., 79, 4269 (1974).

J. P. McClure in Equatorial Anomaly, F. DeMendonca ed. Brazil Space Commission (1965).

PUBLICATIONS

Atmospheric conditions in 1995 ..., J. Atmos. Sol. Terr. Phys. (in Press).